

CUIDADO É FUNDAMENTAL

Escola de Enfermagem Alfredo Pinto – UNIRIO

INTEGRATIVE REVIEW OF LITERATURE

DOI: 10.9789/2175-5361.rpcfo.v16.13122

NURSING CARE FOR USERS DIAGNOSED WITH CROHN'S DISEASE: AN INTEGRATIVE REVIEW

*Cuidados de enfermagem ao usuário diagnosticado com doença de crohn: uma revisão integrativa**Cuidados de enfermería a usuarios diagnosticados de enfermedad de Crohn: una revisión integradora***Maria Alice Nascimento de Lima¹** **José Nacélio da Silva Ferreira²** **Andréa Couto Feitosa³** **Janayle Kéllen Duarte de Sales⁴** **Ana Maria Machado Borges⁵** **Hercules Pereira Coelho⁶** 

ABSTRACT

Objective: to identify the nursing care provided to patients diagnosed with Crohn's disease. **Method:** this is an integrative literature review carried out in the Latin American and Caribbean Health Sciences Literature, Nursing Database, Medical Literature Analysis and Retrieval System Online, Scientific Electronic Library Online and Google Scholar. The sample consisted of seven studies. **Results:** the analysis of the studies reveals a series of significant results in the context of nursing care, including: the need to plan nursing care adapted to the particularities of the disease; the growing relevance of telehealth; symptom monitoring, nutritional guidance, medication administration and emotional support, emphasizing the individualization of this care. **Final considerations:** the main results of the study highlight the importance of nursing care centered on the user's well-being, disease control, emotional support and encouraging self-care.

DESCRIPTORS: Crohn's disease; Regional enteritis; nursing Care;

^{1,2,3,5,6}Dr. Leão Sampaio University Center (UNILEÃO). Ceará, Juazeiro do Norte, Brazil.

⁴Regional University of Cariri (URCA). Ceará, Crato, Brazil.

Received: 19/02/2024; Accepted: 04/04/2024; Published online: 12/05/2024

Corresponding Author: José Nacélio da Silva Ferreira nacelliosilva@gmail.com

How cited: Lima MAN, Ferreira JNS, Feitosa AC, Sales JKD, Borges AMM, Coelho HP. Nursing care for users diagnosed with crohn's disease: an integrative review. *R Pesq Cuid Fundam* [Internet]. 2023 [cited year month day];16:e13122. Available from:

<https://doi.org/10.9789/2175-5361.rpcfo.v16.13122>



RESUMO

Objetivo: identificar os cuidados de enfermagem direcionados a usuários diagnosticados com a doença de Crohn. **Método:** trata-se de uma revisão integrativa da literatura, realizada na Literatura Latino-Americana e do Caribe em Ciências da Saúde, Base de Dados em Enfermagem, *Medical Literature Analysis and Retrieval System Online*, *Scientific Electronic Library Online* e *Google Acadêmico*. A amostra foi composta por sete estudos. **Resultados:** a análise dos estudos revela uma série de resultados significativos no contexto dos cuidados de enfermagem, podendo citar: a necessidade de planejar a assistência de enfermagem adaptada às particularidades da doença; a crescente relevância da telessaúde; monitoramento de sintomas, orientação nutricional, administração de medicamentos e apoio emocional, enfatizando a individualização desses cuidados. **Considerações finais:** os principais resultados do estudo destacam a importância dos cuidados de enfermagem centrados no bem-estar do usuário, controle da doença, apoio emocional e estímulo ao autocuidado.

DESCRIPTORS: Doença de crohn; Enterite regional; Cuidados de enfermagem;

RESUMEN

Objetivos: identificar los cuidados de enfermería prestados a pacientes con diagnóstico de enfermedad de Crohn. **Método:** se trata de una revisión bibliográfica integradora realizada en la Literatura Latinoamericana y del Caribe en Ciencias de la Salud, Base de Datos de Enfermería, *Medical Literature Analysis and Retrieval System Online*, *Scientific Electronic Library Online* y *Google Scholar*. La muestra estuvo constituida por siete estudios. **Resultados:** el análisis de los estudios revela una serie de resultados significativos en el contexto de los cuidados de enfermería, entre ellos: la necesidad de planificar cuidados de enfermería adaptados a las particularidades de la enfermedad; la creciente relevancia de la telesalud; la monitorización de síntomas, la orientación nutricional, la administración de medicamentos y el apoyo emocional, destacando la individualización de estos cuidados. **Consideraciones finales:** los principales resultados del estudio destacan la importancia de los cuidados de enfermería centrados en el bienestar del usuario, el control de la enfermedad, el apoyo emocional y el fomento del autocuidado.

DESCRIPTORES: Enfermedad de crohn; Enteritis regional; Cuidados de enfermería.

INTRODUCTION

According to the Brazilian Association of Ulcerative Colitis and Crohn's Disease (ABCD), Crohn's Disease (CD) is a chronic inflammatory disease of the Gastrointestinal Tract (GIT), which predominantly affects the lower part of the small intestine (ileum) and the large intestine (colon), and can affect any part of the GIT.¹

In Brazil, according to the Brazilian Society of Coloproctology (SBCP), the occurrence of new cases of CD is represented by 7 per 100,000 inhabitants, with a higher concentration mainly in the Southeast and South.² The conditions that predispose to the development of CD are related to immunological and environmental factors, such as: sedentary lifestyle, poor diet, alcoholism and smoking, which are directly related to the risk of CD.³

The main symptoms of CD mentioned in the scientific literature are: abdominal cramps, diarrhoea (which may or may not be associated with haematochezia), vomiting, fever and weight loss. It can also cause complications outside the GIT, such as skin rashes, arthritis and inflammation of the eyes.⁴⁻⁵

During the progression of CD, one or more complications can occur, namely: ulcers, fistulas, abdominal abscesses, intestinal obstructions caused by thickening of the wall of the affected site, malnutrition and cholelithiasis, due to poor absorption of certain substances.⁶ Other, less frequent complications are cancer of the large intestine and digestive haemorrhages.⁷⁻⁸⁻⁹

CD can affect practically all systems and organs, either locally or systemically. Extra-intestinal manifestations can

accompany or appear after the onset of intestinal alterations, which can be joint, eye, dermatological, hepatobiliary, nephrological, haematological, vascular, pancreatic, pulmonary and cardiac alterations.¹⁰

As it is a chronic disease, treatment for CD aims to control symptoms, delay the progression of the disease and improve the quality of life of users. Anti-inflammatory and immunosuppressive medications are used to treat this comorbidity, with the aim of relieving symptoms, prolonging the remission of the disease and postponing possible surgical interventions.¹¹

In the meantime, nurses are part of this context, guiding and facilitating care for individuals living with CD, and they must provide qualified and equitable assistance. To this end, nurses must make use of available health technologies, which can help to achieve systematised, organised and resolute nursing care.

The main nursing care provided to users with CD is related to education and health promotion for self-care and lifestyle changes; guidance on diet, maintaining self-esteem and the use of non-steroidal anti-inflammatory drugs (NSAIDs); encouraging the construction of coping strategies, among others.¹²

This study is justified by the need to understand the main nursing care provided to patients diagnosed with CD, with a view to identifying the risks and implications of the disease, as well as the difficulties faced by patients living with this pathology, in order to understand the paths to equitable care.

The aim of this study was therefore to identify the nursing care provided to patients diagnosed with Crohn's disease.

METHOD

This is a descriptive integrative literature review, carried out in accordance with the following stages: development of the guiding question; literature search or sampling; data collection; critical evaluation of the studies included in the review; discussion of the review's results; and presentation of the integrative review method.¹³

The Population, Variables and Outcomes (PVO) strategy was used to draw up the guiding research question, as shown in Chart 1.

Chart 1 - Elaboration of the guiding question using the PVO strategy. Juazeiro do Norte, CE, Brazil, 2023.

Strategy Items	Components	DeCS	MeSH
Population	Usuários com doença de Crohn e/ou Enterite regional	Doença de Crohn OR Enterite Regional	<i>Crohn Disease</i>
Variables	Assistência de enfermagem	Cuidados de Enfermagem	<i>Nursing Care</i>
Outcomes	Melhoria dos sinais e sintomas	Qualidade de vida	<i>Quality of Life</i>

DeCS: *Descritores em Ciências da Saúde*; MeSH: *Medical Subject Headings*; OR: *Ou*.

Source: Data extracted from the study (own elaboration).

After applying the PVO strategy, the guiding question of the research was: what nursing care is provided to patients diagnosed with Crohn's disease?

The search and selection of studies was carried out in several databases, between July and August 2023, in a paired manner, including the Latin American and Caribbean Literature in Health Sciences (LILACS) and the Nursing Database (BDENF), via the Virtual Health Library (VHL); the Medical Literature Analysis and Retrieval System Online (MEDLINE), via PubMed; the academic search engine Google Scholar; as well as the journal directory Scientific Electronic Library Online (SciELO).

The Boolean operators AND and OR were used to cross-reference the Health Sciences Descriptors (DeCS) and their corresponding Medical Subject Headings (MeSH). Chart 2 shows the search strategies derived from the various DeCS/MeSH combinations used in the study.

Inclusion criteria were: primary scientific articles published in full, free of charge in journals available in the selected databases; and articles published between 2017 and 2022 (the last six years).

The period from 2017 to 2022 was chosen to ensure the inclusion of the most recent evidence on nursing care for users diagnosed with CD. This ensures that the review reflects contemporary practices, updates to guidelines and

Chart 2 - Search strategy for articles by crossing DeCS and MeSH. Juazeiro do Norte, CE, Brazil, 2023.

Databases	Search strategies (DeCS and MeSH)
LILACS e BDENF	(Doença de Crohn OR Enterite Regional) AND (Cuidados de Enfermagem) AND (Qualidade de Vida); (Doença de Crohn OR Enterite Regional) AND (Cuidados de Enfermagem); (Doença de Crohn OR Enterite Regional) AND (Qualidade de Vida).
SciELO e MEDLINE	(<i>Crohn Disease</i>) AND (<i>Nursing Care</i>) AND (<i>Quality of Life</i>); (<i>Crohn Disease</i>) AND (<i>Nursing Care</i>); (<i>Crohn Disease</i>) AND (<i>Quality of Life</i>).
Google Acadêmico	("Doença de Crohn" OR "Enterite Regional") AND ("Cuidados de Enfermagem") AND ("Qualidade de vida").

DeCS: *Descritores em Ciências da Saúde*; MeSH: *Medical Subject Headings*; LILACS: *Literatura Latino Americana e do Caribe em Ciências da Saúde*; BDENF: *Base de Dados em Enfermagem*; SciELO: *Scientific Eletronic Library Online*; MEDLINE: *Medical Literature Analysis and Retrieval System Online*; AND: *E*; OR: *Ou*.

Source: Data extracted from the study (own elaboration).

possible innovations that have occurred in the last six years. It is important to emphasise that language was not used as an inclusion criterion in order to avoid possible restrictions on the sample that could introduce linguistic bias into the research.

On the other hand, the exclusion criteria adopted included: duplicate articles in the databases; and articles that did not fit the theme or did not provide answers to the guiding question of the research, by fully reading the titles and abstracts of the articles.

The Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) was used to conduct the search and select the studies, where applicable, as shown in Figure 1.


During the search and selection of studies, 1849 articles were initially identified. During the identification stage, 1696 (91.7%) articles were excluded due to unavailability in full (paid-for studies) or duplication, resulting in 153 (8.3%) studies. After applying the inclusion criteria, 93 (60.8%) records were excluded, leaving 60 (39.2%) articles. When eligibility was analysed, 53 (88.3%) studies were excluded because they addressed another population or did not answer the research question, by reading the full title and abstract. Therefore, the final sample consisted of seven studies (11.7%), indexed in BDENF (28.6%), LILACS (14.3%), MEDLINE (42.86%) and Google Scholar (14.3%).

To analyse the studies, a database was created using Microsoft Office Word (version 2019), which served as the basis

for filing, synthesising, coding and categorising the articles selected for the integrative review.

In consideration of the ethical and legal precepts, it is important to note that this study was not submitted to the Research Ethics Committee (REC), since its methodological nature exempts it from ethical evaluation, in accordance with the guidelines established in Resolutions 466/12 and 510/16.15-16 However, it is essential to emphasise that all the works and sources used to construct this review were duly cited and referenced, following the principles of authorship and copyright.

Figure 1 - Flowchart for identifying, selecting and including studies, according to the recommendations of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA). Juazeiro do Norte, CE, Brazil, 2023.



LILACS: *Literatura Latino Americana e do Caribe em Ciências da Saúde*; BDENF: *Base de Dados em Enfermagem*; SciELO: *Scientific Electronic Library Online*; MEDLINE: *Medical Literature Analysis and Retrieval System Online*.

Source: Page et al., 2020 (Adapted).¹⁴

RESULTS

After carrying out the article search and selection strategy, identification, selection and inclusion, a total of seven studies were identified which, in short, synthesised the main findings regarding nursing care for users diagnosed with CD. Chart 3 provides crucial information on each study.

Analysing the characterisation of the studies, we observed that the highest concentration of publications occurred in the years 2018 and 2021, with two studies published in each year, totalling 57.1% of the articles included in the research.

The publications covered a wide range of journals, both national and international. With regard to the origin of the studies, we observed that the majority of publications originated in Brazil, three studies (42.9%). In addition, the other studies were distributed in different countries, with one study coming from Belgium (14.3%), one from the United Kingdom (14.3%), one from Lebanon (14.3%), and one from Spain (14.3%).

With regard to the methodological approach, the majority of the studies, i.e. four (57.1%), used quantitative methods in their research. As far as NEC is concerned, the vast majority of studies are classified at level 4, six articles (85.7 per cent).

Table 4 provides a concise summary of the main information from each study.

Analysing the seven studies on CD reveals a series of significant results in the context of nursing care. Machado *et al.*¹⁷ emphasise the need to plan nursing care adapted to the particularities of adolescents with the disease. From a contemporary perspective, Navarro-Correal *et al.*¹⁸ emphasise the growing relevance of telehealth, where the "Advanced IBD Nurse (EADII)" plays a crucial role in providing remote care.

In addition, Guimarães; Gonçalves and Silva¹⁹ highlight the different facets of nursing care, including symptom monitoring, nutritional guidance, medication administration and emotional support, emphasising the individualisation of this care. Lopes *et al.*¹² present a comprehensive list of nursing activities for users with CD.

Chart 3 - Characterisation of the articles included in the integrative review. Juazeiro do Norte, CE, Brazil, 2023.

A	Article title	Authors, year and origin	Magazine / Journal and Database	Approach	NEC [†]
1	Aspectos sociodemográficos e clínicos relacionados à doença de crohn em adolescentes	Machado <i>et al.</i> , 2021 (Brasil) ¹⁷	Enferm Foco (BDENF)	Estudo Quantitativo	4
2	Impact of the COVID-19 pandemic on the activity of advanced-practice nurses on a reference unit for inflammatory bowel disease	Navarro-Correal <i>et al.</i> , 2021 (Espanha) ¹⁸	Gastroenterología y Hepatología (MEDLINE)	Estudo Qualitativo	4
3	Doença de Crohn: um estudo de caso	Guimarães; Gonçalves; Silva, 2020 (Brasil) ¹⁹	FINOM (Google Acadêmico)	Estudo de caso	5

4	<i>Diagnosis and nursing activities for the care of patient with Crohn's disease</i>	Lopes et al., 2019 (Brasil) ¹²	Rev Enferm UFPI (BDENF)	Estudo Quantitativo	4
5	<i>Refocusing IBD patient management: personalized, proactive, and patient-centered care</i>	Siegel, 2018 (Líbano) ²⁰	<i>The American Journal of gastroenterology</i> (LILACS)	Estudo Quantitativo	4
6	<i>Second N-ECCO consensus statements on the european nursing roles in caring for patients with Crohn's Disease or Ulcerative Colitis</i>	Kemp et al., 2018 (Reino Unido) ²¹	<i>Journal of Crohn's and Colitis</i> (MEDLINE)	Estudo Quantitativo	4
7	<i>Effects of introduction of an inflammatory bowel disease nurse position on the quality of delivered care</i>	Coenen et al., 2017 (Bélgica) ²²	<i>European Journal of Gastroenterology & Hepatology</i> (MEDLINE)	Estudo Qualitativo	4

†Souza; Silva; Carvalho (2010).

A: Codificação do artigo; NEC: Nivel de Evidência Científica; LILACS: Literatura Latino Americana e do Caribe em Ciências da Saúde; BDENF: Base de Dados em Enfermagem; SciELO: Scientific Eletronic Library Online; MEDLINE: Medical Literature Analysis and Retrievel System Online.

Source: Data extracted from the study (own elaboration).

Chart 4 - Synthesis of the results of the studies included in the integrative review. Juazeiro do Norte, CE, Brazil, 2023.

A	Authors and year	Aim of the study	Main results
1	Machado et al., 2021 ¹⁷	Descrever as características sociodemográficas e clínicas de adolescentes com doença de Crohn.	Although it does not indicate specific care, the study highlights the importance of evaluating the sociodemographic and clinical characteristics of Crohn's Disease in adolescents, in order to provide nursing/health care planning based on the peculiarities of the health-disease process, with an impact on improving the health condition and well-being of users with Crohn's Disease.
2	Navarro-Correal et al., 2021 ¹⁸	Relatar o impacto da pandemia de COVID-19 na atividade dos enfermeiros que trabalham numa unidade de Doença Inflamatória Intestinal (DII) e identificar as razões para a prestação de cuidados de saúde à distância e a sua relação com determinadas características.	The study highlights that the Crohn's-Colitis Care Unit (UACC) of a University Hospital in Barcelona follows a model of comprehensive patient-centred care through face-to-face and distance care, with emphasis on the role of the Advanced Inflammatory Bowel Disease (IBD) Nurse in Telehealth Assistance (TA), known as telemedicine, due to their clinical profile and professional skills. It is possible that these professionals could provide more detailed information on specific nursing care for users with IBD.
3	Guimarães; Gonçalves; Silva, 2020 ¹⁹	Realizar um estudo de caso, evidenciando o papel do enfermeiro no cuidado e acompanhamento do paciente portador da Doença de Crohn.	Nursing care aimed at users diagnosed with Crohn's disease includes: symptom monitoring, user education, nutritional guidance, medication administration, pain management, prevention and treatment of complications, emotional support, and self-care education. It is important to stress that nursing care must be individualised, taking into account the specific needs of each user. In addition, collaboration with other health professionals, such as doctors, nutritionists and psychologists, is essential for comprehensive and effective care.

4	Lopes <i>et al.</i> , 2019 ¹²	Elaborar e validar diagnósticos e atividades de enfermagem para cuidado ao paciente com Crohn.	This study provides a list of 62 nursing activities aimed at users with Crohn's disease, which were drawn up on the basis of 26 clinically validated nursing diagnoses. These activities include, for example, monitoring vital signs, administering medication, nutritional counselling, education about the disease and its symptoms, and emotional support.
5	Siegel, 2018 ²⁰	Identificar os pacientes com alto risco de ter complicações da sua doença, para que possamos interromper esta história natural e alterar o seu curso.	According to the study, nurses' attention should be centred on the individual patient and what we can do for them now, i.e. provide them with an optimal quality of life and prevent complications related to Inflammatory Bowel Disease (IBD) in the future.
6	Kemp <i>et al.</i> , 2018 ²¹	Melhorar os cuidados prestados aos doentes com Doença Inflamatória Intestinal (DII) através do desenvolvimento de diretrizes, educação e investigação.	The study addresses the fact that Inflammatory Bowel Disease (IBD) is highly complex and requires specialised nursing care and management. To inform, inspire and improve standards of care in IBD by providing evidence-based guidelines that enable IBD nurses to move from fundamental to advanced care, encouraging research by IBD nurses. With this, new and expanding IBD services will also serve as a basis for creating educational programmes for Inflammatory Bowel Disease nurses.
7	Coenen <i>et al.</i> , 2017 ²²	Tornar possível que o enfermeiro responsável pela doença inflamatória intestinal (DII) seja o principal ponto de acesso para educação, aconselhamento e apoio. Investigando o efeito da introdução de um enfermeiro de Doença Inflamatória Intestinal na qualidade dos cuidados prestados.	Nursing care for patients diagnosed with Crohn's disease includes long-term outpatient follow-up, maintenance therapy and acute interventions for disease flares. The Inflammatory Bowel Disease (IBD) nurse plays an important role in this care, providing counselling, support and education to users and their families so that they understand the pathophysiology and follow-up strategies in IBD. The introduction of an IBD nurse in a tertiary referral centre has proved effective in reducing emergency admissions and unscheduled outpatient visits, improving the quality of care and reducing costs.

A: Codificação do artigo. DII: Doença Inflamatória Intestinal; ET AL: E Outros; DE: Diagnóstico de Enfermagem; TA: Assistência à Telessaúde; UACC: Unidade de Cuidados de Crohn-Colite.

Source: Data extracted from the study (own elaboration).

Siegel²⁰, in turn, emphasises the importance of improving Quality of Life (QoL) and preventing complications related to IBD. Kemp *et al.*²¹ highlight the complexity of IBD and the need for guidelines and specialised education for nurses. Finally, Coenen *et al.*²² highlight the effectiveness of introducing IBD nurses in improving the quality of care and reducing costs, with an emphasis on education, counselling and support as key components of nursing care.

DISCUSSION

The results underline the vital role of nursing in caring for users with CD, emphasising assessment, personalised care, telehealth and interdisciplinary collaboration in improving users' QoL. That said, analysing the results revealed two categories related to nursing care for users diagnosed with CD, as shown below:

Approaches to nursing care in Crohn's disease

This category discusses the importance of nursing care for users with CD, highlighting the planning of more effective care, the relevance of sociodemographic assessment, symptom monitoring, nutritional guidance, medication administration and mental support.

As explored, nursing offers various care approaches for users diagnosed with CD. It is crucial to stress that nursing care must be individualised, meeting the specific needs of each user. In this context, collaboration with other professionals broadens the scope of treatment and increases its effectiveness.¹⁹

The clinically validated nursing activities and diagnoses, according to the taxonomies, meet the needs of patients diagnosed with CD in outpatient clinics, favouring the provision of resolute nursing care. In general, nursing activities aim to provide a professional service adapted to the user's real health needs, promoting QoL in various dimensions. It can be seen that the main nursing care is centred on QoL, including encouraging the user, raising self-esteem and managing symptoms, among others.¹²

In order to optimise treatment, reduce the number of patients and achieve disease remission, it is essential to focus on the individual needs of each user and take immediate action to improve their QoL and prevent future complications or changes in the course of the disease.²⁰

The inclusion of nurses specialising in IBD in the multidisciplinary team has made them responsible for counselling, support and care, facilitating rapid intervention during disease flares. This has resulted in a significant reduction in emergency consultations and unscheduled outpatient visits, thanks to the effectiveness of nurse management.²²

Considering the active role of nursing in the care process, it is essential that it encourages users diagnosed with CD to adopt self-care practices, providing relevant information on adherence to treatment, diet and mental health, as well as surgical procedures and clinical follow-up.²³

In their study, Silva *et al.*²⁴ emphasise that CD is chronic, requiring prolonged treatment, affecting both the physical and psychosocial health of users, with the potential to change their routines, QoL and well-being. They conclude that social factors have a direct impact on the psychological well-being of users diagnosed with CD.

In line with this, the results of Martinelli's study²⁵ show an improvement in the QoL of individuals with CD through the Systematisation of Nursing Care (SNC). This process proves essential in life, treatment and daily routine, allowing users to achieve a healthier life in their health-disease process, adjusting their plans and dreams to the restrictions and complications of this comorbidity. It is clear to users diagnosed with CD that, with proper nursing care, it is possible to guarantee a satisfactory and lasting degree of QoL, even considering that the disease has no cure.

The impact of telehealth and the role of nurses in Crohn's disease

This section highlights the crucial impact of telehealth and the adaptation of nurses to this new reality in CD care. Thus, this category addresses the influence of telehealth and nurses on various aspects, including accessibility, quality of care and the financial impact on CD management.

Telehealth has emerged as a crucial tool in the provision of remote care. Its relevance in CD care is undeniable, because through this technology, users can receive medical and nursing care flexibly, regardless of location. The flexibility offered by telehealth empowers users, allowing them greater autonomy over their health. In addition, telehealth eliminates geographical barriers, enabling distant users to access specialists and specialised care, contributing to accessibility.¹⁸

Robust evidence has shown that the use of telehealth offers a series of additional benefits. These include a significant reduction in the time it takes to provide care, since users can be seen promptly in their own homes. This approach also contributes to a reduction in costs related to travelling for users and health professionals, relieving the financial burden and stress associated with frequent trips.²⁶

One notable aspect is the improvement in the quality of care provided by telehealth, allowing access to specialists even in remote areas where the presence of highly specialised professionals can be limited. Telehealth's ability to effectively connect users and healthcare professionals, regardless of their geographical location, enhances treatment and support options, resulting in more efficient and effective care.²⁷

Navarro-Correal *et al.*¹⁸ emphasised in their study the relevance of telehealth in CD care and the role of the EADII in providing remote care. Nurses, including EADIIs, play a crucial role in CD care through telehealth. They provide support, education and ongoing monitoring to users, even at a distance. This transition requires a change in practices and the development of new skills to meet users' needs remotely. The ability to adapt to this new type of care is fundamental for effective care and for promoting quality care.¹⁸⁻²⁸

Corroborating this, Santos²⁹, in his dissertation on telecare in the treatment of CD, identified positive results in relation to the reduction in the incidence rate of drug complications related to CD, by monitoring users with this technology.

Thus, users diagnosed with CD can access specialists for their conditions more conveniently, avoiding unnecessary travelling. However, as noted in the study³⁰, remote visits may not be suitable for all user profiles. Therefore, telehealth must be implemented in a personalised way, taking users' individual needs into account to guarantee the quality of care offered.

In addition to the benefits for users and healthcare professionals, telehealth also has financial implications for healthcare management. It can reduce costs by avoiding hospitalisations and frequent travel, providing significant relief for users and the healthcare system. However, as highlighted by Shah *et al.*³¹, the success

of telehealth can be influenced by factors such as the existence of well-defined clinical protocols and accessibility challenges. It is therefore essential to address financial issues carefully and promote equal access to telehealth to ensure that all users benefit from it.

FINAL CONSIDERATIONS

The main results of the study highlight the importance of nursing care centred on the user's well-being, disease control, emotional support and encouraging self-care. Nursing plays a fundamental role in promoting quality of life and in the effective management of Crohn's disease, also showing a significant increase in the integration of telehealth into care.

These results have significant implications for nursing practice, highlighting the need for nursing professionals to seek specialisation and training in order to offer more comprehensive and targeted care to users with Crohn's disease. With the global growth in the prevalence of Crohn's disease, it is essential that nurses are prepared to face specific challenges, including the clinical management of the condition, psychological support and the promotion of self-care.

The study faced some limitations, such as the scarcity of national research on the subject, which restricted the contextualisation of nursing care to the Brazilian reality. In addition, the presence of outdated articles also affected the breadth of the analysis.

Given these limitations, it is suggested that future studies address these knowledge gaps. It is recommended that national research be carried out to explore the main nursing care for users with Crohn's disease in the Brazilian context. It is also crucial to assess the understanding and perspective of users and their families in relation to nursing care, with a view to promoting autonomy in self-care.

REFERENCES

1. Associação Brasileira de Colite Ulcerativa e Doença de Crohn (ABCD). Sobre a doença de crohn: o que é? [Internet]. 2023 [acesso em 29 de abril 2023]. Disponível em: <https://www.abcd.org.br/sobre-a-doenca-de-crohn/>
2. Gasparini RG, Sasaki LY, Saad-Hossne R. Inflammatory bowel disease epidemiology in São Paulo State, Brazil. *Clinical and Experimental Gastroenterology*. [Internet]. 2018 [cited 2023 jul 23];18(11). Available from: <https://doi.org/10.2147/CEG.S176583>
3. Tonsic BR, Cardoso BM. Papel dos imunomoduladores na doença de crohn. In: IX Mostra Interna de Trabalhos de Iniciação Científica [evento na internet]. 2018 out 23-26; Paraná, Brasil [acesso em 17 de maio de 2023]. Disponível em: https://rdu.unicesumar.edu.br/bitstream/123456789/2239/1/barbara_ribeiro_tonsic.pdf
4. Matsuoka K, Kobayashi T, Ueno F, Matsui T, Hirai F, Inoue N, et al. Evidence-based clinical practice guidelines for inflammatory bowel disease. *J. gastroenterol*. [Internet]. 2018 [cited 2023 abr 29];53(3). Available from: <https://doi.org/10.1007/s00535-018-1439-1>
5. Seyedian SS, Nokhostin F, Malamir MD. A review of the diagnosis, prevention, and treatment methods of inflammatory bowel disease. *J. Med Life*. [Internet]. 2019 [cited 2023 mar 29];12(2). Available from: <https://doi.org/10.25122/jml-2018-0075>
6. Venito LS, Santos MSB, Ferraz AR. Doença de Crohn e retocolite ulcerativa. *Rev Eletrônica Acervo Saúde*. [Internet]. 2022 [acesso em 28 de abril de 2023];15(7). Disponível em: <https://doi.org/10.25248/reas.e10667.2022>
7. Mota ES, Kiss DR, Teixeira M G, Almeida MG, Sanfront FA, Habr-Gama A, Cecconello I. Manifestações extra-intestinais em doença de Crohn e retocolite ulcerativa: prevalência e correlação com o diagnóstico, extensão, atividade, tempo de evolução da doença. *Rev bras Coloproct*. [Internet]. 2007 [acesso em 12 de maio de 2023];27(4). Disponível em: <https://doi.org/10.1590/S0101-98802007000400001>
8. World Gastroenterology Organization (WGO). Doença inflamatória intestinal: uma perspectiva global. In: *Practice Guidelines: Manejo clínico da DII*. [Internet]. 2009 [cited 2023 mar 31]. Available from: <https://www.worldgastroenterology.org/UserFiles/file/guidelines/inflammatory-bowel-disease-portuguese-2009.pdf>
9. Araújo SEA, Dias AR, Seid VE, Campos FG, Nahas SC. Videocirurgia no manejo da doença de Crohn intestinal. *Rev bras Coloproct*. [Internet]. 2010 [acesso em 28 de abril de 2023];30(3). Disponível em: <https://doi.org/10.1590/S0101-98802010000300001>
10. Oliveira JA, Viebig RF, Nacif M, Bazanelli AP. Relação do consumo alimentar com sinais e sintomas na doença de Crohn. *Brazilian J. Development*. [Internet]. 2021 [acesso em 26 de abril de 2023];7(7). Disponível em: <https://doi.org/10.34117/bjdv7n7-549>
11. Papacosta NG, Nunes GM, Pacheco RJ, Cardoso MV, Guedes VR. Doença de crohn: um artigo de revisão. *Rev de Patologia do Tocantins*. [Internet]. 2017 [acesso em 01 de abril de 2023];4(2). Disponível em: <https://doi.org/10.20873/uft.2446-6492.2017v4n2p25>
12. Lopes AM, Costa APC, Silva GRF, Fernandes ADBF. Diagnosis and nursing activities for the care of patient with Crohn's disease. *Rev Enferm UFPI*. [Internet]. 2019 [cited 2023 may 10];8(2). Available from: <https://doi.org/10.26694/2238-7234.8245-51>
13. Souza MT, Silva MD, Carvalho R. Revisão integrativa: o que é e como fazer. *Einstein*. [Internet]. 2010 [acesso em 15 de fevereiro de 2023];8(1). Disponível em: <https://doi.org/10.1590/s1679-45082010rw1134>

14. Page MJ, Mckenzie J, Bossuyt P, Boutron I, Hoffmann T, Mulrow CD. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ*. [Internet]. 2021 2019 [cited 2023 aug 11];372(71). Available from: <https://doi.org/10.1136/bmj.n71>
15. Ministério da Saúde (BR). Conselho Nacional de Saúde. Resolução no 466, de 12 de dezembro de 2012. 2023 [Internet]. Brasília: Ministério da Saúde [acesso em 18 de março de 2023]. Disponível em: <http://www.conselho.saude.gov.br/resolucoes/2012/Reso466.pdf>
16. Ministério da Saúde (BR). Conselho Nacional de Saúde. Resolução no 510, de 07 de abril de 2016. 2016 [Internet]. Brasília: Ministério da Saúde [acesso em 19 de março de 2023]. Disponível em: https://www.in.gov.br/materia/asset_publisher/Kujrw0TZC2Mb/content/id/22917581
17. Machado K, Andrade PCST, Toledo TT, Peres EM, Gomes HF, Almeida IS, et al. Aspectos sociodemográficos e clínicos relacionados à doença de crohn em adolescentes. *Enferm Foco*. [Internet]. 2021 [acesso em 10 de maio de 2023]; 2(5). Disponível em: <https://doi.org/10.21675/2357-707X.2021.v12.n5.4523>
18. Navarro-Correal S, Borrueal N, Robles V, Guise CH, Ayala LFM, Martinez ZP, et al. Impact of the COVID-19 pandemic on the activity of advanced-practice nurses on a reference unit for inflammatory bowel disease. *Gastroenterología y Hepatología*. [Internet]. 2019 [cited 2023 aug 15];44(7). Available from: <https://doi.org/10.1016/j.gastrohep.2020.11.018>
19. Guimarães MC, Gonçalves MDS, Silva CP. Doença de Crohn: um estudo de caso. *Humanidades e Tecnologia em Revista (Finom)*. [Internet]. 2020 [acesso em 30 de abril de 2023];23(1). Disponível em: http://revistas.icesp.br/index.php/FINOM_Humanidade_Tecnologia/article/view/1187/865
20. Siegel CA. Refocusing IBD Patient Management: Personalized, Proactive, and Patient-Centered Care. *The American J. Gastroenterology*. [Internet]. 2018 [cited 2023 aug 19];113(10). Available from: <https://doi.org/10.1038/s41395-018-0246-x>
21. Kemp K, Dibley L, Chauhan U, Greveson K, Jäghult S, Ashton K, et al. Second N-ECCO Consensus Statements on the European Nursing Roles in Caring for Patients with Crohn's Disease or Ulcerative Colitis. *J. Crohn's and Colitis*. [Internet]. 2018 [cited 2023 may 04];12(7). Available from: <https://doi.org/10.1093/ecco-jcc/jjy020>
22. Coenen S, Weyts E, Vermiere S, Ferrante M, Noman M, Ballet V, et al. Effects of introduction of an inflammatory bowel disease nurse position on the quality of delivered care. *European J. Gastroenterology & Hepatology*. [Internet]. 2017 [cited 2023 may 11];29(6). Available from: <https://doi.org/10.1097/MEG.0000000000000839>
23. Organização Brasileira de Doença de Crohn e Colite (GEBIIB). Os cuidados de Enfermagem em Doença Inflamatória Intestinal [Internet]. 2019 [acesso em 21 de junho de 2023]. Disponível em: <https://gediib.org.br/pg/noticias-pg/os-cuidados-de-enfermagem-em-doenca-inflamatoria-intestinal/>
24. Silva DM, Mello JM, Schneider LCL, Santana DMG. Papel da enterografia no diagnóstico da Doença de Crohn. *Arquivos do MUDI*. [Internet]. 2019 [acesso em 12 maio de 2023];23(3). Disponível em: <https://periodicos.uem.br/ojs/index.php/ArqMudi/article/download/51517/751375149128/>.
25. Martinelli CL. Análise da qualidade de vida de um indivíduo portador da doença de Crohn após prescrição de enfermagem [Graduação em Enfermagem]. Espírito Santo (Brasil): Católica de Vitória Centro Universitário; 2018. [acesso em 03 de agosto de 2023]. Disponível em: https://unisaes.br/wp-content/uploads/2021/10/2018-1_Enf_Charlene.pdf
26. Caetano R, Silva AB, Guedes ACCM, Paiva CCN, Ribeiro GR, Santos DL, Silva RM. Desafios e oportunidades para telessaúde em tempos da pandemia pela COVID-19: uma reflexão sobre os espaços e iniciativas no contexto brasileiro. *Cad. Saúde Pública*. [Internet]. 2020 [acesso em 12 de maio de 2023];36(5). Disponível em: <https://doi.org/10.1590/0102-311X00088920>
27. Portnoy J, Waller M, Elliott T. Telemedicine in the Era of COVID-19. *J Allergy Clin Immunol Pract*. [Internet]. 2020 [cited 2023 may 12];8(5). Available from: <https://doi.org/10.1016/j.jaip.2020.03.008>
28. Barbosa IA, Silva MJP. Nursing care by telehealth: what is the influence of distance on communication?. *Rev Bras Enferm*. [Internet]. 2017 [cited 2023 may 25];70(5). Available from: <http://dx.doi.org/10.1590/0034-7167-2016-0142>
29. Santos RM. O telecuidado no tratamento das doenças inflamatórias intestinais: ensaio clínico randomizado. [Mestrado Profissional em Enfermagem Assistencial]. Rio de Janeiro (Brasil): Universidade Federal Fluminense; 2016. [acesso em 01 de novembro de 2023]. Disponível em: <https://app.uff.br/riuff/bitstream/handle/1/5894/Rachael%20Miranda%20dos%20Santos.pdf?sequence=1&isAllowed=y>
30. Hoyo J, Millán M, Garrido-Marim A, Nos P, Acosta MB, Bujanda L, et al. Changes in the management of IBD patients since the onset of COVID-19 pandemic. A path toward the implementation of telemedicine in Spain?. *Gastroenterol Hepatol*. [Internet]. 2022 [cited 2023 may 12];45(9). Available from: <https://doi.org/10.1016/j.gastrohep.2021.08.006>
31. Shah KP, Triana AJ, Gusdorf RE, McCoy AB, Pabla B, Scoville E, Dalal R, et al. Demographic Factors Associated With Successful Telehealth Visits in Inflammatory Bowel Disease Patients. *Inflamm Bowel Dis*. [Internet]. 2022 [cited 2023 apr 22];28(3). Available from: <https://doi.org/10.1093/ibd/izab068>.